



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE EXPANSION OF $\text{sn } x$.

By MR. ROLLIN A. HARRIS, Jamestown, N. Y.

(1.) Since $\text{sn}(-x) = -\text{sn } x$, we may assume

$$\begin{aligned}\text{sn } x &= A_0x + A_1x^3 + A_2x^5 + \dots; \\ \dots \text{sn } y &= A_0y + A_1y^3 + A_2y^5 + \dots, \\ \text{sn}(x+y) &= A_0(x+y) + A_1(x+y)^3 + \dots, \\ \text{cn } x \text{ dn } x &= \text{sn}'x = A_0 + 3A_1x^2 + 5A_2x^4 + \dots;\end{aligned}$$

and so for $\text{cn } y \text{ dn } y$.

Substituting these developments in the formula

$$\text{sn}(x+y) = \frac{\text{sn}^2x - \text{sn}^2y}{\text{sn } x \text{ cn } y \text{ dn } y - \text{sn } y \text{ cn } x \text{ dn } x},$$

we have, after clearing of fractions,

$$\begin{aligned}& [A_0(x+y) + A_1(x+y)^3 + \dots] \\ & \times \left[\begin{aligned} & (A_0x + A_1x^3 + \dots)(A_0 + 3A_1y^2 + \dots) \\ & - (A_0y + A_1y^3 + \dots)(A_0 + 3A_1x^2 + \dots) \end{aligned} \right] \\ & = (A_0x + A_1x^3 + \dots)^2 - (A_0y + A_1y^3 + \dots)^2. \quad (a)\end{aligned}$$

Since the second member contains terms of the form cx^n or cy^n only, it follows that, if the first member be expanded and similar terms united, the coefficient of any resulting term, whose form is cx^py^q where $p, q > 0$, must be equal to zero. We could thus obtain relations between the coefficients $A_0, A_1, \dots A_n$. But actual multiplication is not necessary, since we are not concerned with the entire product. We proceed as follows:—

(2.) Performing the multiplication indicated in the first term of the second factor of the first member of (a), we have

$$\begin{aligned}(A_0x + A_1x^3 + \dots)(A_0 + 3A_1y^2 + \dots) \\ = A_0A_0y^0x + A_0A_1y^0x^3 + A_0A_2y^0x^5 + \dots \\ + 3A_1A_0y^2x + 3A_1A_1y^2x^3 + 3A_1A_2y^2x^5 + \dots \\ + 5A_2A_0y^4x + 5A_2A_1y^4x^3 + 5A_2A_2y^4x^5 + \dots \\ \dots \dots \dots\end{aligned}$$

		Order in x .				
		1	3	5	7	
Order in y .	0	$A_0 A_0$	$A_0 A_1$	$A_0 A_2$	$A_0 A_3$. . .
	2	$3A_1 A_0$	$3A_1 A_1$	$3A_1 A_2$	$3A_1 A_3$. . .
	4	$5A_2 A_0$	$5A_2 A_1$	$5A_2 A_2$	$5A_2 A_3$. . .
	6	$7A_3 A_0$	$7A_3 A_1$	$7A_3 A_2$	$7A_3 A_3$. . .
	

(β)

(3.) Because of symmetry, if x and y of this table be interchanged, we have the product of $A_0 y + A_1 y^3 + \dots$ and $A_0 + 3A_1 x^2 + \dots$

(4.) The first factor of (α) may be written $\sum_{n=0}^{\infty} A_n (x+y)^{2n+1}$,

	0	1	2	3	4	5	6	7
0		A_0		A_1		A_2		A_3
1	A_0		$3A_1$		$5A_2$		$7A_3$	
2		$3A_1$		$10A_2$		$21A_3$		
3	A_1		$10A_2$		$35A_3$			
4		$5A_2$. . .				
5	A_2		. . .					
6		. . .						
7	. . .							

(γ)

or

Suppose that we multiply (β) by (γ) and wish to find all terms in which x and y have the exponents p and q , respectively. Each of such terms will consist of two factors $c_{st} x^s y^t$, $c_{uv} x^u y^v$, the one from (β), the other from (γ), where $s + u = p$ and $t + v = q$. Since $s \leq p$ and $t \leq q$, we see at once, what factors can come from (β); and since $p = s + u$ and $q = t + v$, the factors from (γ) are determined. In fact, we have only to multiply the first term of (β) by that term of (γ) whose order

is the highest permissible; again, the next higher in (β) with reference to either variable by the next lower in (γ) with reference to the same variable, and so continue until s and t have reached their limits.

(5.) EXAMPLE. Let it be required to find the entire coefficient of $x^5 y^3$.

From what has just been said, (β) multiplied by (γ) would give as the coefficient of $x^5 y^3$

$$A_0 A_0 \cdot 35 A_3 + A_0 A_1 \cdot 10 A_2 + A_0 A_2 \cdot A_1 \\ + 3 A_1 A_0 \cdot 5 A_2 + 3 A_1 A_1 \cdot 3 A_1 + 3 A_1 A_2 \cdot A_0.$$

Now interchange x and y ; there results as the coefficient of $y^5 x^3$

$$A_0 A_0 \cdot 21 A_3 + A_0 A_1 \cdot A_2 \\ + 3 A_1 A_0 \cdot 10 A_2 + 3 A_1 A_1 \cdot A_1 \\ + 5 A_2 A_0 \cdot 3 A_1 + 5 A_2 A_1 \cdot A_0.$$

But these must be equal, since the entire coefficient of $x^5 y^3$ equals zero [(3), (4)]. Collecting and transposing, we have

$$7 A_3 A_0^2 - 11 A_2 A_1 A_0 + 3 A_1^3 = 0.$$

From other considerations, we know that $A_0 = 1$;

$$\therefore 7 A_3 - 11 A_2 A_1 + 3 A_1^3 = 0.$$

(6.) From (γ) , it is evident that we can always bring A_m into an equation involving A 's of lower subscript only, if we take $p + q = 2(m + 1)$. For example, we could have obtained the above equation by using $x^6 y^2$ instead of $x^5 y^3$. Obviously, the smaller we can take q , the more simple will be the operation.

If $q = 1$, the result is an identity, and so of no value to us; but if $q = 2$, the result in general will not be an identity. We observe that the coefficient of the A_m , $q = 2$, from (γ) is the same as the coefficient of the third term of $(a + b)^{2m+1}$, or $m(2m + 1)$. In (β) , A_m does not occur while we are considering $x^p y^2$. Now interchange x and y ; i. e. consider $x^2 y^p$. The A_m from (γ) has the same coefficient as the second term of $(a + b)^{2m+1}$, or $2m + 1$; from (β) , it also has the coefficient $2m + 1$. These added together give $2(2m + 1)$; but $2(2m + 1)$ is not equal to $m(2m + 1)$, except for $m = 2$. Since, in general, we can obtain the required relations by making $q = 2$, we shall hereafter confine ourselves to that value. The operation then becomes very simple. E.g. Required the value of A_4 .

Here $p + q = 10$; therefore $p = 8$. We have, then,

$$A_0(A_0 \cdot 36 A_4 + A_1 \cdot 21 A_3 + A_2 \cdot 10 A_2 + A_3 \cdot 3 A_1) \\ + 3 A_1(A_0 \cdot A_3 + A_1 \cdot A_2 + A_2 \cdot A_1 + A_3 \cdot A_0) \\ \Rightarrow A_0(A_0 \cdot 9 A_4 + 3 A_1 \cdot 7 A_3 + 5 A_2 \cdot 5 A_2 + 7 A_3 \cdot 3 A_1 + 9 A_4 \cdot A_0); \\ \therefore 6 A_4 - 4 A_3 A_1 - 5 A_2^2 + 2 A_2 A_1^2 = 0.$$

(7.) The relation sought is embodied in the following general equation:—

$$\begin{aligned}
 & A_0 \sum_{r=1}^{r=m} r(2r+1) A_r A_{m-r} + 3A_1 \sum_{r=1}^{r=m} A_{r-1} A_{m-r} \\
 &= A_0 \sum_{r=1}^{r=m+1} (2r-1)(2m-2r+3) A_{r-1} A_{m-r+1}; \\
 & \left\{ A_0 \sum_{r=2}^{r=m+1} (2r-1)(2m-2r+3) A_{r-1} A_{m-r+1} \right. \\
 & \quad \left. - 3A_1 \sum_{r=1}^{r=m} A_{r-1} A_{m-r} - A_0 \sum_{r=1}^{r=m-1} r(2r+1) A_r A_{m-r} \right\} \\
 \therefore A_m &= \frac{(m-2)(2m+1) A_0^2}{(m-2)(2m+1) A_0^2} \\
 m \text{ odd} \\
 A_m &= \frac{\left\{ 2A_0 \sum_{r=2}^{r=\frac{1}{2}(m+1)} (2r-1)(2m-2r+3) A_{r-1} A_{m-r+1} \right. \\
 & \quad \left. - 6A_1 \sum_{r=1}^{r=\frac{1}{2}(m-1)} A_{r-1} A_{m-r} - 3A_1 A_{\frac{1}{2}(m-1)}^2 - \sum_{r=1}^{r=m-1} r(2r+1) A_r A_{m-r} \right\}}{(m-2)(2m+1) A_0^2} \\
 m \text{ even} \\
 A_m &= \frac{\left\{ 2A_0 \sum_{r=2}^{r=\frac{1}{2}m} (2r-1)(2m-2r+3) A_{r-1} A_{m-r+1} + (m+1)^2 A_0 A_{\frac{1}{2}m}^2 \right. \\
 & \quad \left. - 6A_1 \sum_{r=1}^{r=\frac{1}{2}m} A_{r-1} A_{m-r} - A_0 \sum_{r=1}^{r=m-1} r(2r+1) A_r A_{m-r} \right\}}{(m-2)(2m+1) A_0^2}
 \end{aligned}$$

By giving m the values 3, 4, . . . , we have

$$\begin{aligned}
 7A_3 - 11A_2A_1 + 3A_1^3 &= 0, \\
 6A_4 - 4A_3A_1 - 5A_2^2 + 2A_2A_1^2 &= 0, \\
 11A_5 - 3A_4A_1 - 13A_3A_2 + 2A_3A_1^2 + A_2^2A_1 &= 0, \\
 26A_6 - A_5A_1 - 22A_4A_2 + 3A_4A_1^2 - 14A_3^2 + 3A_3A_2A_1 &= 0. \\
 &\dots\dots\dots
 \end{aligned}$$

OBSERVATIONS.

All terms are of order three, counting the A_0 's, and the weight of each term of any of these equations equals the greatest subscript, equals the number of terms.

A_1 and A_2 are supposed to be known; they may be called the *source* of all succeeding ones. For in terms of these two, any that follow may be expressed.

NOTE.—A similar method is applicable to $\cos x$ and $\operatorname{dn} x$.